

10. Jahangiri M, Kreutzer J, Zurakowski D, Bacha E, Jonas RA. Evaluation of hemostatic and coagulation factor abnormalities in patients undergoing the Fontan operation. *J Thorac Cardiovasc Surg.* 2000; 120:778-82.
11. Odegard KC, McGowan FX, DiNardo JA, et al. Coagulation factor abnormalities in patients with single ventricle physiology precede the Fontan operation. *J Thorac Cardiovasc Surg.* 2002;123:459-65.
12. Odegard KC, McGowan FX, Zurakowski D, et al. Coagulation factor abnormalities in patients with single ventricle physiology immediately prior to the Fontan procedure. *Ann Thorac Surg.* 2002;73:1770-7.
13. Andrew M, Vegh P, Johnston M, Bowker J, Ofosu F, Mitchell L. Maturation of the hemostatic system during childhood. *Blood.* 1992; 80:1998-2005.
14. Rauch R, Ries M, Hofbeck M, Buheitel G, Singer H, Klinge J. Hemostatic changes following the modified Fontan operation (total cavopulmonary connection). *Thromb Haemost.* 2000;83:678-82.
15. Kyle PA, Minar E, Hirsch M, et al. High Plasma levels of factor VIII and the risk of recurrent venous thromboembolism. *N Engl J Med.* 2000;343:457-62.
16. O'Donnell J, Tuddenham EG, Manning R, Kemball-Cook G, Johnson D, Laffan M. High prevalence of elevated factor VIII levels in patients referred for thrombophilia screening: role of increased synthesis and relationship to the acute phase reaction. *Thromb Haemost.* 1997;77: 825-8.
17. Kamphuisen PW, Eikenboom JC, Vos HL, et al. Increased levels of factor VIII and fibrinogen in patients with venous thrombosis are not caused by acute phase reaction. *Thromb Haemost.* 1999;81:680-3.
18. Kraaijenhagen RA, in't Anker PS, Koopman MM, et al. High plasma concentration of factor VIIIc is a major risk factor for venous thrombembolism. *Thromb Haemost.* 2000;83:5-9.
19. Kamphuisen PW, Eikenboom PW, Bertina RM. Elevated factor VIII levels and the risk of thrombosis. *Arterioscler Thromb Vasc Biol.* 2001;21:731-8.
20. Koster T, Blann AD, Briet E, Vandenbroucke JP, Rosendaal FR. Role of clotting factor VIII in effect of von Willebrand factor on occurrence of deep-vein thrombosis. *Lancet.* 1995;345:152-5.
21. Hollestelle MJ, Thinnis T, Crain K, et al. Tissue distribution of factor VIII gene expression in vivo—a closer look. *Thromb Haemost.* 2001; 86:855-61.

ON THE MOVE?

Send us your new address at least six weeks ahead

Don't miss a single issue of the journal! To ensure prompt service when you change your address, please photocopy and complete the form below.

Please send your change of address notification at least six weeks before your move to ensure continued service. We regret we cannot guarantee replacement of issues missed due to late notification.

JOURNAL TITLE:

Fill in the title of the journal here. _____

OLD ADDRESS:

Affix the address label from a recent issue of the journal here.

NEW ADDRESS:

Clearly print your new address here.

Name _____

Address _____

City/State/ZIP _____

COPY AND MAIL THIS FORM TO:

Mosby
Subscription Customer Service
6277 Sea Harbor Dr
Orlando, FL 32887

OR FAX TO:

407-363-9661



OR PHONE:

800-654-2452
Outside the U.S., call
407-345-4000